



ONLY USED FOR ANIMAL HEALTH

**POXDOLL**

INJECTABLE SUSPENSION LYOPHILIZATE

**LYOPHILIZED SHEEP-GOATS POX VACCINE**

**NAME AND BRIEF DESCRIPTION OF THE VETERINARY BIOLOGICAL PRODUCT:**

POXDOLL is a lyophilized, live, attenuated vaccine used against pox disease in sheep-goats and Lumpy Skin Disease in cattle.

**Name and power of the active substances, and auxiliary substances, if it exist:**

**COMPOSITION:** The following minimum ratios are used for each dose.

Components	Amount	Power
Attenuated SPV (Bk) strain	$10^{2.5}$ TCID <sub>50</sub> /dose	Immunizing antigen
Lactalbumin hydrolysate	0,25 mg / dose	Preservative
Sucrose	0,50 mg / dose	Preservative

**INDICATIONS:**

POXDOLL is used for the active immunization of sheep and goat against sheep & goat pox, as well cattle against Lumpy Skin disease.

**CONTRAINDICATIONS:**

There are no contraindications.

**ADVERSE EFFECTS:**

In some animals, there may be hard or soft swellings of 0.2-5 cm in diameter and a temporary rise in body temperature in the vaccine administration area.

**TARGET SPECIES:**

Sheep, goats and cattle

**METHOD OF ADMINISTRATION AND DOSE:**

0.2 ml for 6-12 week old lambs and kids, 0.5 ml for animals older than 12 weeks old and 3 ml for cattle older than 6 months (subcutaneous).

**WITHDRAWAL PERIOD OF ANIMAL ORIGIN PRODUCTS:**

None.

**PRESENTATION:**

50 or 100 doses lyophilized vaccine in Glass vials, each 20 vials are packaged in plastic viol. The corresponding diluent in 25 and 50 ml brown glass bottles unboxed or boxed in Styrofoam 100 bottles each.

ONLY USED FOR ANIMAL HEALTH

**POXDOLL**

INJECTABLE SUSPENSION LYOPHILIZATE



**LYOPHILIZED SHEEP-GOATS POX VACCINE**

**STORAGE CONDITIONS AND SHELF LIFE :**

Real-time shelf life is 36 months at 2 / 8°C in the dark.

**WARNINGS :**

Infected animals or animals within incubation period should not be vaccinated. The diluted vaccine should be used within two hours at the latest and shaken well before use. The vaccine is not administered to pregnant sheep and goats in the last 1/3 of pregnancy (in the last 6 weeks of pregnancy), in the first month following birth, in places where the disease does not occur, and to kids and lambs under 6 weeks of age. Calves born from vaccinated animals should not be vaccinated before 6 months of age. Calves born from unvaccinated animals should be vaccinated before 6 months of age. Safety in pregnant cattle has not been proven precisely, so it is recommended not to use it in the last 3 weeks of pregnancy.

**PRECAUTIONS FOR THE PERSON ADMINISTERING THE VACCINE AND RECOMMENDATIONS FOR VETERINARIANS:**

The lyophilized pellet should be homogenized after dissolving it thoroughly with the vaccine diluent using sterile injectors. During the vaccine administration, care should be taken not to infect the eyes or hands, and if any contamination occurs, the area should be cleaned with plenty of water and disinfected.

**DISPOSAL OF WASTE AND UNUSED PRODUCTS:**

Vaccines that are not opened and used are destroyed by boiling for 10 minutes. Unopened and unused vaccines should be sent to the manufacturer for disposal.

“Keep out of reach of children”

“Consult veterinarian if any undesired effects occur”



**THE MARKETING AUTHORIZATION ISSUED BY MINISTRY OF AGRICULTURE AND FORESTRY, DATE AND NUMBER :** 25.06.2007

**NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER:**

Dollvet Veteriner Aşı İlaç Biyolojik Madde Üretim Sanayi ve Ticaret A.Ş.  
Mehmet Nesih Özmen Mah. Kasım Sok. D Blok Apt. No:7 Güngören/İSTANBUL - TURKEY  
**E-mail:** dollvet@dollvet.com.tr / **Phone:** +90 212 422 02 01 / **Fax:** +90 212 422 00 72

**NAME AND ADDRESS OF THE PLACE OF PRODUCTION:**

Dollvet Veteriner Aşı İlaç Biyolojik Madde Üretim Sanayi ve Ticaret A.Ş.  
Organize Sanayi Bölgesi Koçören OSB Mahallesi 106. Cadde No:4 Eyyübiye/SANLIURFA - TURKEY  
**E-mail:** dollvet@dollvet.com.tr / **Phone:** +90 414 369 11 33 / **Fax:** +90 414 369 16 62