

FOR VETERINARY USE ONLY

LSD-NDOLL

LIVE, ATTENUATED AND LYOPHILIZED VACCINE AGAINST
NODULAR EXANTHEMA (LUMPY SKIN) DISEASE IN CATTLE



Dollvet

LYOPHILISATE FOR SUSPENSION FOR INJECTION AND SOLVENT

COMPOSITION:

Following minimum quantities are used in each dose for each dosage (2 ml),

Ingredients	Quantity	Function
Attenuated LSD Neethling strain	10^{3-5} TCID ₅₀ /dose	Immunizing antigen
Lactalbumin hydrolysate	0,25 mg/dose	Preservative
Sucrose	0,50 mg/dose	

INDICATIONS:

LSD-NDOLL is used against nodular exanthema (Lumpy Skin Disease) in cattle, and produces immunity by generating a cellular and humoral immune response in target animals 21 days after vaccination.

CONTRAINDICATIONS:

None.

SIDE EFFECTS:

In some cases, hard or soft swellings 0,2-5 cm in diameter at the injection site and a temporary rise in body temperature may be observed.

DOSAGE AND ADMINISTRATION ROUTE:

Dose for cattle is 2 ml, administered by subcutaneous injection.

WITHDRAWAL PERIODS:

None.

PRESENTATION:

For vaccine lyophilization: 5 and 10 ml Type I neutral glass vials, with aluminum caps and butyl stoppers, containing respectively 10, 25, 50 and 100 cattle doses, For solvent: 20, 50, 100 and 200 ml Type II neutral glass or plastic bottles, closed by butyl stoppers and aluminum caps are used.

STORAGE CONDITIONS AND SHELF LIFE:

Lyophilized vaccine stored at + 2/8°C, and in the dark. Shelf life is 2 years. Do not freeze.

Solvent should be store at room temperature below + 25°C. The solvent should be cooled to + 2/8°C before use.

FOR VETERINARY USE ONLY

LSD-NDOLL

LIVE, ATTENUATED AND LYOPHILIZED VACCINE AGAINST
NODULAR EXANTHEMA (LUMPY SKIN) DISEASE IN CATTLE



Dollvet

LYOPHILISATE FOR SUSPENSION FOR INJECTION AND SOLVENT

WARNINGS:

Vaccines should be stored at +2/+8°C in a dark place and shaken well before each use. Opened vials should be used within 2 hours, and necessary precautions should be taken to ensure vaccines are transferred at storage temperature.

PRECAUTIONS TO BE TAKEN BY THE PERSON ADMINISTERING THE VACCINE AND RECOMMENDATIONS FOR VETERINARIANS:

Using sterile syringes, the lyophilized pellet should be homogenized, thoroughly dissolving it with the solvent. During vaccination, special care should be taken not to contaminate the eyes or hands. In case of an accidental contact, the site should be cleaned with water and medical attention should be sought if needed.

DISPOSAL OF WASTE AND UNUSED PRODUCTS:

Unused or opened but not completely used vaccines are disposed of as medical waste, according to the provisions of the applicable waste regulations.

GENERAL WARNINGS:

Keep out of reach of children.

Consult a veterinarian upon an unexpected effect.

LEAFLET APPROVAL DATE: 23.10.2018

DATE OF MARKETING AUTHORIZATION ISSUED BY MINISTRY OF AGRICULTURE AND FORESTRY: 23.10.2018

NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER:

Dollvet Biyoteknoloji A.Ş.

Konaklar Mah. Akasyalı Sok. No:10

Beşiktaş /İSTANBUL

E-mail: dollvet@dollvet.com.tr Phone: +90 212 422 02 01 Fax: +90 212 422 00 72

NAME AND ADDRESS OF PRODUCTION SITE:

Dollvet Biyoteknoloji A.Ş.

Organize Sanayi Bölgesi Koçören OSB Mahallesi 106. Cadde No:6

Eyyübiye/ŞANLIURFA

E-mail: dollvet@dollvet.com.tr Phone: +90 414 369 11 33 Fax: +90 414 369 16 62

200360/R.01