

FOR VETERINARY USE ONLY

IMMUDOLL^{CG}

NON-SPECIFIC IMMUNOSTIMULANT



Dollvet

SUSPENSION FOR INJECTION

COMPOSITION:

Active ingredients	
Inactivated <i>Cutibacterium granulorum</i>	20 mg/ml
Excipients	
Ethanol	12,5%

IMMUNOLOGICAL PROPERTIES:

Pharmacotherapeutic group: Other immunologics (Non-specific immunostimulant)

ATCvet code: QI01AX, QI02AX, QI03AX, QI04AX

This biological product stimulates a non-specific immune response. By inducing the reticulo-endothelial system, it activates non-specific cell response (monocytes, macrophages, neutrophils and cytotoxic cells), provides inflammation and stimulates interferon production.

INDICATIONS:

It is used to increase non-specific immunity in target species. It increases the organism's resistance against infections. It helps treatment by strengthening the effectiveness of therapeutic intervention in diseases. It contributes to the development of the general immune response by increasing the non-specific immune response. It increases body resistance in situations that cause stress (transport or environmental conditions).

METHOD OF ADMINISTRATION AND DOSAGE:

In cattle, sheep and goats, 0.2 ml/10 kg body weight should be administered subcutaneously or intramuscularly. In chickens, intramuscular administration should be done at a dose of 0.05 ml/animal per animal.

In cattle, sheep and goats:

For preventive purposes, 2 doses are recommended with 7 days intervals. In pregnant animals, 2 doses are recommended with 7 days intervals, 30-60 days before birth. As an aid for treatment, it is recommended to apply 2 doses with an interval of 3 days.

Chickens:

5 ml of the product is diluted with 50 ml of an aseptic solvent or an aseptic isotonic solvent and administered intramuscularly as 0.5 ml per animal.

SPECIAL CLINICAL INFORMATION AND SPECIAL WARNINGS FOR TARGET SPECIES:

IMMUDOLL-CG should be shaken well before use.

Bottles should be brought to room temperature before application.

Aseptic needles and syringes should be used in every application.

Use During Pregnancy and Lactation

It is safe to use in cattle, sheep and goats during the last period of pregnancy and lactation.

ADVERSE EFFECTS:

Short-term local reactions (redness, slight swelling and crusting) may occur at the injection site. In very rare cases, anaphylactic reactions may occur in sensitive animals. In this case, symptomatic treatment (antihistamine/antiallergic drugs) should be applied.

INTERACTION WITH OTHER MEDICINAL PRODUCTS:

IMMUDOLL-CG stimulates non-specific immunity, so when used in combination with other immunological products, it is expected to support the acquired immunity. There is no information about the compatibility of this product when used with other drugs.

SYMPTOMS, PRECAUTIONS AND ANTIDOTE IN CASE OF OVERDOSE:

No overdose studies have been conducted as it is an inactivated vaccine.

FOR VETERINARY USE ONLY

IMMUDOLL CG

NON-SPECIFIC IMMUNOSTIMULANT



Dollvet

SUSPENSION FOR INJECTION

WITHDRAWAL PERIOD:

Withdrawal period is 0 day.

CONTRAINDICATIONS:

It has no known contraindications.

GENERAL WARNINGS:

“Keep out of reach of children”

“Consult your veterinarian before use and in case of an unexpected effect”

PRECAUTIONS FOR PRACTITIONERS AND RECOMMENDATIONS FOR VETERINARIANS:

The product has no known negative effects on human health, general asepsis and antiseptics rules must be followed. In case of injuries that may occur during the application of the biological product, the area should be cleaned and disinfected.

The vaccine should only be administered by a veterinarian or a Veterinary health technician under the supervision of a veterinarian.

STORAGE CONDITIONS AND SHELF LIFE:

Store at +2/+8°C and in the dark. Do not freeze.

Shelf life is 24 months. Opened vials should be used within 6-8 hours at 20-25 °C.

DISPOSAL OF WASTE AND WARNINGS FOR NON-TARGET SPECIES:

Unused immunostimulant biological products and immunostimulant biological products opened for use are disposed of as medical waste according to the current waste regulations.

PRESENTATION:

IMMUDOLL-CG is available in 5 ml amber Type I glass vials, in 25, 50 and 100 ml amber Type II glass vials or in 50 and 100 ml white polypropylene bottles. Glass and polypropylene bottles sealed with grey bromobutyl rubber stopper and aluminum cap.

IMMUDOLL-CG individually packed in plastic vials or cardboard boxes: 1x5 ml, 10x5 ml, 1x25 ml, 1x50 ml and 1x100 ml. Not all packaging forms may be marketed.

LEAFLET APPROVAL DATE: 05.02.2024

DATE OF MARKETING AUTHORIZATION ISSUED BY THE MINISTRY OF AGRICULTURE AND FORESTRY: 05.02.2024

NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER:

Dollvet Biyoteknoloji Anonim Şirketi

Konaklar Mah. Akasyalı Sok. No:10 Beşiktaş /İSTANBUL

Phone: +90 212 422 02 01 Fax: +90 212 422 00 72 E-mail: dollvet@dollvet.com.tr

NAME AND ADDRESS OF MANUFACTURER:

Dollvet Biyoteknoloji Anonim Şirketi

Organize Sanayi Bölgesi Koçören OSB Mahallesi 106. Cadde No:6 Eyyübiye/ŞANLIURFA

Phone: +90 414 369 11 33 Fax: +90 414 369 16 62 E-mail: dollvet@dollvet.com.tr

201006-R.00